

Integral inequalities for generalized approximately h -convex functions on fractal sets via generalized local fractional integrals

Research Article

Maria Bibi^{1*} and Muhammad Muddassar^{1†}

1 Department of Basics Sciences, University of Engineering and Technology, Taxila. Pakistan

Abstract: The present paper study the notion of generalized approximately h -convex function and define a novel generalized fractional integral operator on fractal sets. We derive local fractional integral inequalities involving newly defined generalized local fractional integral operator for generalized approximately h -convex function. Special cases of the results are also established.

MSC: 26D15, 26A33, 26A51, 26D10, 90C23

Keywords: Generalized approximately h -convex functions • Local fractional integral inequality • Fractal Sets • Local Fractional Integrals

Received 2022-06-02; Accepted 2022-07-09; Published 2022-07-14

1. Introduction

The study of convex functions offered provoking results in multiple areas of mathematics e.g, mathematical economics, linear programming, optimization, dynamic systems and control theory. Nowadays, different kinds of integral inequalities concerning convex functions and their generalizations are getting attention of researchers to work on. More information can be found in the references [3–7, 15, 16, 19].

Definition 1.1.

A mapping $F : I \subseteq R \rightarrow R$ is known to be convex, if

$$F(\ell r + (1 - \ell)s) \leq \ell F(r) + (1 - \ell)F(s) \quad (1)$$

If $\forall r, s \in I, \ell \in [0, 1]$

* Corresponding author

* E-mail: mariabibi782@gmail.com

† E-mail: muhammad.mudassar@uettaxila.edu.pk

Famous inequality derived by J.Hadamard [9] in 1881 is known as Hermite-Hadamard Inequality. Let $F : I \subseteq R \rightarrow R$, be a convex function if $r, s \in I$ and $r < s$, then

$$F\left(\frac{r+s}{2}\right) \leq \frac{1}{s-r} \int_r^s F(y)dy \leq \frac{F(r)+F(s)}{2} \quad (2)$$

S. Varosanec [26] presented the notion of h -convex functions.

Definition 1.2.

Let $h : J \subseteq R$ be non negative function and $h \neq 0$. A non negative function $F : I = [r, s] \rightarrow R$ is known to be h -convex, if for all $\ell \in [0, 1]$.

$$F(\ell r + (1-\ell)s) \leq h(\ell)F(r) + h(1-\ell)F(s) \quad (3)$$

In [12] Kashuri et al defined approximately h -convex function. Consider $(X, ||, ||_I)$ be a normed quasi linear space, and I is nonempty convex subset of $X, d : X \times X \rightarrow R$.

Definition 1.3.

Consider $h : [0, 1] \rightarrow R$ be non negative function and $h \neq 0$. A function $F : I \rightarrow R$ is called to be approximately h -convex, if

$$F(\ell r + (1-\ell)s) \leq h(\ell)F(r) + h(1-\ell)F(s) + d(r, s) \quad (4)$$

holds for all $\ell \in (0, 1)$ and $r, s \in I$

Sarikaya and Ertuğral in [17] presented the left-sided and right-sided generalized fractional integral operators, as:

$${}_{r+}I_{\varphi}F() = \int_r^{\varphi(-\ell)} \frac{\varphi(-\ell)}{(-\ell)} F(\ell)d\ell, > r \quad (5)$$

$${}_{s-}I_{\varphi}F() = \int_{\varphi(\ell-)}^s \frac{\varphi(\ell-)}{(\ell-)} F(\ell)d\ell, < s \quad (6)$$

Hadamard inequality for the generalized fractional integral operators is established by Sarikaya and Ertuğral in [17].

Theorem 1.1.

Let $F : [r, s] \rightarrow R$ be a convex function on $[r, s]$, with $r < s$, then for fractional integral operators the following inequality holds:

$$F\left(\frac{r+s}{2}\right) \leq \frac{1}{2\Lambda(1)} [{}_{r+}I_{\varphi}F(s) + {}_{s-}I_{\varphi}F(r)] \leq \frac{F(r)+F(s)}{2} \quad (7)$$

where the mapping $\Lambda(x) : [0, 1] \rightarrow R$ is given by

$$\Lambda() := \int_0^1 \frac{\varphi((s-r)\ell)}{\ell} d\ell \quad (8)$$

Kashuri et al. [11], established inequalities of Hermite Hadamard type for approximately h -convex functions involving generalized fractional integrals.

Theorem 1.2.

Let $F : [r, s] \rightarrow R$ be an approximately h -convex on $[r, s]$, with $r < s$. Then for generalized fractional integrals the following inequality holds:

$$\begin{aligned} & \frac{1}{2h(\frac{1}{2})} F\left(\frac{r+s}{2}\right) - A_1 \leq \frac{1}{2\Lambda(1)} [{}_r I_\varphi F(s) + {}_s I_\varphi F(r)] \\ & \leq \frac{[F(r) + F(s)]}{2\Lambda(1)} \int_0^1 \frac{\varphi((s-r)\ell)}{\ell} [h(\ell) + h(1-\ell)] d\ell + d(r, s) \end{aligned}$$

where,

$$A_1 := \frac{1}{2\Lambda(1)h(\frac{1}{2})} \int_r^s \frac{\varphi(s-)}{s-} d(r+s-).$$

We now discuss some preliminaries of local fractional calculus theory introduced by Yang in [28, 29]. These concepts and consequences are linked with fractal order derivatives and integrals.

If $r_1^\varsigma, r_2^\varsigma, r_3^\varsigma \in R^\varsigma$ ($0 < \varsigma \leq 1$), then

- $r_1^\varsigma + r_2^\varsigma \in R^\varsigma, r_1^\varsigma r_2^\varsigma \in R^\varsigma,$
- $r_1^\varsigma + r_2^\varsigma = r_2^\varsigma + r_1^\varsigma = (r_1 + r_2)^\varsigma = (r_2 + r_1)^\varsigma,$
- $r_1^\varsigma + (r_2^\varsigma + r_3^\varsigma) = (r_1 + r_2)^\varsigma + r_3^\varsigma,$
- $r_1^\varsigma r_2^\varsigma = r_2^\varsigma r_1^\varsigma = (r_1 r_2)^\varsigma = (r_2 r_1)^\varsigma,$
- $r_1^\varsigma (r_2^\varsigma r_3^\varsigma) = (r_1^\varsigma r_2^\varsigma) r_3^\varsigma,$
- $r_1^\varsigma (r_2^\varsigma + r_3^\varsigma) = r_1^\varsigma r_2^\varsigma + r_1^\varsigma r_3^\varsigma,$
- $r_1^\varsigma + 0^\varsigma = 0^\varsigma + r_1^\varsigma = r_1^\varsigma,$ and $r_1^\varsigma 1^\varsigma = 1^\varsigma r_1^\varsigma = r_1^\varsigma$
- If $r_1^\varsigma < r_2^\varsigma,$ then $r_1^\varsigma + r_3^\varsigma < r_2^\varsigma + r_3^\varsigma,$
- If $0^\varsigma < r_1^\varsigma, 0^\varsigma < r_2^\varsigma,$ then $0^\varsigma < r_1^\varsigma r_2^\varsigma,$

Local fractional derivative and integral on R^ς are defined as,

Definition 1.4.

([28, 29]) A non-differentiable function $F : R \rightarrow R^\varsigma, y \rightarrow F(y)$ is local fractional continuous at $y_0,$ if for any $\epsilon > 0,$ there exists $\delta > 0$ such that

$$|F(y) - F(y_0)| < \epsilon^\varsigma$$

holds $|y - y_0| < \delta,$ with $\epsilon, \delta \in R.$ If $F(y)$ is local continuous on $(c, d),$ and we denote $F(y) \in C_\varsigma(c, d).$

Definition 1.5.

([28, 29]) Local fractional derivative of the function $F(y)$ of order ς at $y = y_0$ can be defined as

$$F^{(\varsigma)}(y_0) = \left. \frac{d^\varsigma F(y)}{d\ell^\varsigma} \right|_{y=y_0} = \lim_{y \rightarrow y_0} \frac{\Gamma(1+\varsigma)(F(y) - F(y_0))}{(y - y_0)}$$

$D_\varsigma(b, c)$ is ς -local derivative set. If there exists $F^{((K+1)\varsigma)}(y) = \overbrace{D_y^\varsigma \dots D_y^\varsigma}^{(n+1)\text{times}} F(y)$ for any $y \in I \subseteq R$, we denote $F \in D_{(n+1)\varsigma}(I)$, and $n = 0, 1, 2, \dots$

Definition 1.6.

([28, 29]) Let $F(y) \in C_\varsigma[c, d]$. Local fractional integral of $F(w)$ can be defined by

$${}_b I_c^\varsigma F(y) = \frac{1}{\Gamma(1+\varsigma)} \int_b^c F(\ell)(d\ell)^\varsigma = \frac{1}{\Gamma(1+\varsigma)} \lim_{\Delta\ell \rightarrow 0} \sum_{f=0}^{N-1} F(\ell_f)(\Delta\ell_f)^\varsigma,$$

Where $c = \ell_0 < \ell_1 < \dots < \ell_{N-1} < \ell_N = d, [\ell_f, \ell_{f+1}]$ is partition of $[c, d]$, $\Delta\ell_f = \ell_{f+1} - \ell_f, \Delta\ell = \max\{\ell_0, \ell_1, \dots, \ell_{N-1}\}$.

Note that ${}_c I_c^\varsigma F(y) = 0$ and ${}_c I_d^\varsigma F(y) = -{}_d I_c^\varsigma F(y)$ if $c < d$. We denote $F(y) \in I_y^\varsigma[c, d]$ if there exists ${}_b I_y^\varsigma F(y)$ for any $y \in [b, c]$.

Lemma 1.1.

([28, 29])

1. Let $g(y) = f^{(\varsigma)}(y) \in C_\varsigma[c, d]$, then

$${}_b I_c^\varsigma g(y) = f(d) - f(c)$$

2. Let $g(y), f(y) \in D_\varsigma[c, d]$ and $g^{(\varsigma)}(y), f^{(\varsigma)}(y) \in C_\varsigma[b, c]$, then

$${}_b I_c^\varsigma g(y) f^{(\varsigma)}(y) = g(y) f(y) \Big|_b^c - {}_b I_c^\varsigma g^{(\varsigma)}(y) f(y).$$

Lemma 1.2.

([28, 29])

$$\begin{aligned} \frac{d^\varsigma y^{s\varsigma}}{du^\varsigma} &= \frac{\Gamma(1+s\varsigma)}{\Gamma(1+(s-1)\varsigma)} y^{(s-1)\varsigma}; \\ \frac{1}{\Gamma(\varsigma+1)} \int_b^c y^{s\varsigma} (du)^\varsigma &= \frac{\Gamma(1+s\varsigma)}{\Gamma(1+(s+1)\varsigma)} (d^{(s+1)\varsigma} - c^{(s+1)\varsigma}), \quad s > 0 \end{aligned}$$

Lemma 1.3.

([28, 29])

$${}_b I_c^\varsigma 1^\varsigma = \frac{(d-c)^\varsigma}{\Gamma(1+\varsigma)}$$

Lemma 1.4.

([28, 29]) (Generalized Hölder's inequality) Let $p, q > 1$ with $p^{-1} + q^{-1} = 1$, let $g(w), f(w) \in C_\varsigma[c, d]$, Then

$$\frac{1}{\Gamma(\varsigma+1)} \int_b^c |g(y)f(y)|(dy)^\varsigma \leq \left(\frac{1}{\Gamma(\varsigma+1)} \int_b^c |g(y)|^p (dy)^\varsigma \right)^{\frac{1}{p}} \left(\frac{1}{\Gamma(\varsigma+1)} \int_b^c |f(y)|^q (dy)^\varsigma \right)^{\frac{1}{q}}$$

Recall generalized beta function:

$$B_{\varsigma}(y, x) = \frac{1}{\Gamma(1 + \varsigma)} \int_0^1 \ell^{(y-1)\varsigma} (1 - \ell)^{(x-1)\varsigma} (d\ell)^{\varsigma}, \quad y > 0, x > 0 \quad (9)$$

Local fractional theory has solid applications in control theory, communication engineering, random walk process and Physics [1, 13, 27, 30]. Many researchers studied various types of integral inequalities for generalized definitions of convexity on fractal sets.[2, 8, 10, 14, 18]. Wenbing Sun established various types of Integral inequalities for different generalizations of convex functions in fractal theory [20–25]. According to our knowledge the study of approximately h -convex functions has not been carried out in fractal domain. In this research paper we study the concept of generalized approximately h -convex functions on fractal sets, we also present a novel fractional integral operator on fractal sets through which we establish generalized local fractional integral inequalities.

2. Main Results

We define a function $\varphi_{\beta} : [0, \infty) \rightarrow [0, \infty)$ satisfying the following condition, if for $X^{\varsigma}, Y^{\varsigma}, Z^{\varsigma} > 0$ and independence of $a^{\varsigma}, b^{\varsigma} > 0$

$$\begin{aligned} & \frac{1}{\Gamma(1 + \varsigma)} \int_0^1 \frac{\varphi_{\varsigma}(\ell^{\varsigma})}{\ell^{\varsigma}} (d\ell)^{\varsigma} < +\infty \\ & \frac{1}{X^{\varsigma}} \leq \frac{\varphi_{\varsigma}(b^{\varsigma})}{\varphi_{\varsigma}(a^{\varsigma})} \leq X^{\varsigma}, \quad \text{for } \frac{1}{2^{\varsigma}} \leq \frac{b^{\varsigma}}{a^{\varsigma}} \leq 2^{\varsigma} \\ & \frac{\varphi_{\varsigma}(a^{\varsigma})}{a^{2\varsigma}} \leq Y^{\varsigma} \frac{\varphi_{\varsigma}(b^{\varsigma})}{b^{2\varsigma}} \quad \text{for } b^{\varsigma} < a^{\varsigma} \\ & \left| \frac{\varphi_{\varsigma}(a^{\varsigma})}{a^{2\varsigma}} - \frac{\varphi_{\varsigma}(b^{\varsigma})}{b^{2\varsigma}} \right| \leq Z^{\varsigma} |a^{\varsigma} - b^{\varsigma}| \frac{\varphi_{\varsigma}(a^{\varsigma})}{a^{2\varsigma}} \quad \text{for } \frac{1}{2^{\varsigma}} \leq \frac{b^{\varsigma}}{a^{\varsigma}} \leq 2^{\varsigma} \end{aligned}$$

Now we define generalized local fractional integral operators,

$${}_{r+}I_{\varphi}^{\varsigma}F() = \frac{1}{\Gamma(1 + \varsigma)} \int_r^{\ell} \frac{\varphi_{\varsigma}(-\ell)^{\varsigma}}{(-\ell)^{\varsigma}} F(\ell)(d\ell)^{\varsigma}, \quad > r \quad (10)$$

$${}_{s-}I_{\varphi}^{\varsigma}F() = \frac{1}{\Gamma(1 + \varsigma)} \int^s \frac{\varphi_{\varsigma}(\ell-)^{\varsigma}}{(\ell-)^{\varsigma}} F(\ell)(d\ell)^{\varsigma}, \quad < s \quad (11)$$

Definition 2.1.

Suppose $h : J \rightarrow R$ be non negative function and $h^{\varsigma} \not\equiv 0$. A function $F : I \rightarrow R^{\varsigma} (0 < \varsigma \leq 1)$ be ς order fractal dimensional. If F be non-negative ($F \geq 0^{\varsigma}$) and for $r, s \in I$ and $\ell \in (0, 1)$, then

$$F(\ell r + (1 - \ell)s) \leq h^{\varsigma}(\ell)F(r) + h^{\varsigma}(1 - \ell)F(s) + d^{\varsigma}(r, s) \quad (12)$$

is called to be generalized approximately h -convex function on fractal sets.

Remark 2.1.

If $h^{\varsigma}(\ell) = \ell^{\varsigma}$, the generalized approximately h -convex function turned into generalized approximately convex function. If $h^{\varsigma}(\ell) = \ell^{\varsigma}(1 - \ell)^{\varsigma}$, $h^{\varsigma}(\ell) = 1$, $h^{\varsigma}(\ell) = \ell^{s\varsigma}$, the generalized approximately h -convex function turned into generalized approximately tgs-convex function, generalized approximately P -convex function, generalized approximately s -convex function respectively.

Theorem 2.1.

Let $F : [r, s] \rightarrow R^\varsigma$ and $F(\cdot) \in I^\varsigma[r, s]$ be an approximately h -convex on $[r, s]$, with $0 \leq r < s$. Then following generalized local fractional integrals inequalities hold:

$$\begin{aligned} & \frac{1}{2^\varsigma h^\varsigma(\frac{1}{2})} F\left(\frac{r+s}{2}\right) - B_1 \leq \frac{1}{2^\varsigma \Lambda^\varsigma(1)} [{}_{r+}I_\varphi^\varsigma F(s) + {}_{s-}I_\varphi^\varsigma F(r)] \\ & \leq \frac{[F(r) + F(s)]}{2^\varsigma \Lambda^\varsigma(1)} \frac{1}{\Gamma(1+\varsigma)} \int_0^1 \frac{\varphi_\varsigma((s-r)\ell)^\varsigma}{\ell^\varsigma} [h^\varsigma(\ell) + h^\varsigma(1-\ell)] (d\ell)^\varsigma + d^\varsigma(r, s) \end{aligned}$$

where,

$$B_1 := \frac{1}{2\Lambda^\varsigma(1)h^\varsigma(\frac{1}{2})\Gamma(1+\varsigma)} \int_r^s \frac{\varphi_\varsigma(s-)^^\varsigma}{(s-)^^\varsigma} d^\varsigma(r, r+s-)(d)^\varsigma.$$

Proof. As F is generalized approximately h -convex function, so we have following inequality

$$F\left(\frac{u+v}{2}\right) \leq h^\varsigma\left(\frac{1}{2}\right)[F(u) + F(v)] + d^\varsigma(u, v) \quad (13)$$

Taking $u = \ell r + (1-\ell)s$ and $v = \ell s + (1-\ell)r$

$$\begin{aligned} & \frac{1}{h^\varsigma(\frac{1}{2})} F\left(\frac{r+s}{2}\right) \leq F(\ell r + (1-\ell)s) \\ & + F(\ell s + (1-\ell)r) + \frac{1}{h^\varsigma(\frac{1}{2})} d(\ell r + (1-\ell)s, \ell s + (1-\ell)r) \end{aligned}$$

Multiply both sides of above inequality by $\frac{\varphi_\varsigma((s-r)\ell)}{\ell}$ and integrate local fractionally with respect to ℓ over $[0, 1]$, we have

$$\begin{aligned} & \frac{1}{h^\varsigma(\frac{1}{2})} F\left(\frac{r+s}{2}\right) \frac{1}{\Gamma(1+\varsigma)} \int_0^1 \frac{\varphi_\varsigma((s-r)\ell)}{\ell} (d\ell)^\varsigma \\ & \leq \frac{1}{\Gamma(1+\varsigma)} \int_0^1 \frac{\varphi_\varsigma((s-r)\ell)}{\ell} F(\ell r + (1-\ell)s) (d\ell)^\varsigma + \\ & \frac{1}{\Gamma(1+\varsigma)} \int_0^1 \frac{\varphi_\varsigma((s-r)\ell)}{\ell} F(\ell s + (1-\ell)r) (d\ell)^\varsigma \\ & + \frac{1}{h^\varsigma(\frac{1}{2})\Gamma(1+\varsigma)} \int_0^1 \frac{\varphi_\varsigma((s-r)\ell)}{\ell} d(\ell r + (1-\ell)s, \ell s + (1-\ell)r) (d\ell)^\varsigma \end{aligned}$$

As a consequence, we get

$$\begin{aligned} & \frac{1}{2^\varsigma h^\varsigma(\frac{1}{2})} F\left(\frac{r+s}{2}\right) \leq \frac{1}{2^\varsigma \Lambda^\varsigma(1)} [{}_{r+}I_\varphi^\varsigma F(s) + {}_{s-}I_\varphi^\varsigma F(r)] \\ & + \frac{1}{2^\varsigma \Lambda^\varsigma(1)h^\varsigma(\frac{1}{2})\Gamma(1+\varsigma)} \int_r^s \frac{\varphi_\varsigma(s-)^^\varsigma}{(s-)^^\varsigma} d^\varsigma(r, r+s-)(d)^\varsigma. \end{aligned}$$

we get the proof of right hand side of the inequality.

To prove left hand side of the inequality, since F is approximately h -convex, we have

$$F(\ell r + (1-\ell)s) + F(\ell s + (1-\ell)r) \leq [F(r) + F(s)][h^\varsigma(\ell) + h^\varsigma(1-\ell)] + 2^\varsigma d^\varsigma(r, s)$$

Multiply both sides of inequality by $\frac{\varphi((s-r)\ell)}{\ell}$ and integrating local fractionally with respect to ℓ over $[0, 1]$, we have,

$$\begin{aligned} & \frac{1}{\Gamma(1+\varsigma)} \int_0^1 \frac{\varphi((s-r)\ell)}{\ell} F(\ell u + (1-\ell)v)(d\ell)^\varsigma + \\ & \frac{1}{\Gamma(1+\varsigma)} \int_0^1 \frac{\varphi((s-r)\ell)}{\ell} F(\ell v + (1-\ell)u)(d\ell)^\varsigma + 2^\varsigma d^\varsigma(r, s) \frac{1}{\Gamma(1+\varsigma)} \int_0^1 \frac{\varphi((s-r)\ell)}{\ell} (d\ell)^\varsigma \\ & \leq [F(r) + F(s)] \frac{1}{\Gamma(1+\varsigma)} \int_0^1 \frac{\varphi((s-r)\ell)}{\ell} [h^\varsigma(\ell) + h^\varsigma(1-\ell)](d\ell)^\varsigma + 2^\varsigma d^\varsigma(r, s) \Lambda^\varsigma(1) \end{aligned}$$

As a result, we get

$$\begin{aligned} & \frac{1}{2^\varsigma \Lambda^\varsigma(1)} [{}_{r+} I_\varphi^\varsigma F(s) + {}_{s-} I_\varphi^\varsigma F(r)] \\ & \leq \frac{[F(r) + F(s)]}{2^\varsigma \Lambda^\varsigma(1) \Gamma(1+\varsigma)} \int_0^1 \frac{\varphi((s-r)\ell)}{\ell} [h^\varsigma(\ell) + h^\varsigma(1-\ell)](d\ell)^\varsigma + d^\varsigma(r, s) \end{aligned}$$

□

Corollary 2.1.

If $\varphi(\ell) = \ell^\varsigma$ then following inequality holds

$$\begin{aligned} & \frac{1}{2^\varsigma h^\varsigma(\frac{1}{2})} F\left(\frac{r+s}{2}\right) - B_1 \leq \frac{1}{2^\varsigma (s-r)^\varsigma} [{}_{r+} I^\varsigma F(s) + {}_{s-} I^\varsigma F(r)] \\ & \leq \frac{[F(r) + F(s)]}{2^\varsigma} \frac{1}{\Gamma(1+\varsigma)} \int_0^1 [h^\varsigma(\ell) + h^\varsigma(1-\ell)](d\ell)^\varsigma + \frac{d^\varsigma(r, s)}{(s-r)^\varsigma} \end{aligned}$$

where,

$$B_1 := \frac{1}{2^\varsigma h^\varsigma(\frac{1}{2}) \Gamma(1+\varsigma) (s-r)^\varsigma} \int_r^s d^\varsigma(r, r+s-)(d)^\varsigma.$$

Corollary 2.2.

If $h(\ell) = \ell^\varsigma$ in above inequality then following inequality holds

$$\begin{aligned} & \frac{1}{2^\varsigma h^\varsigma(\frac{1}{2})} F\left(\frac{r+s}{2}\right) - B_1 \leq \frac{1}{2^\varsigma (s-r)^\varsigma} [{}_{r+} I^\varsigma F(s) + {}_{s-} I^\varsigma F(r)] \\ & \leq [F(r) + F(s)] \frac{\Gamma(1+\varsigma)}{\Gamma(1+2\varsigma)} + \frac{d^\varsigma(r, s)}{(s-r)^\varsigma} \end{aligned}$$

Corollary 2.3.

If $h(\ell) = \ell^\varsigma(1-\ell)^\varsigma$ in above inequality then following inequality holds

$$\begin{aligned} & \frac{1}{2^\varsigma h^\varsigma(\frac{1}{2})} F\left(\frac{r+s}{2}\right) - B_1 \leq \frac{1}{2^\varsigma (s-r)^\varsigma} [{}_{r+} I^\varsigma F(s) + {}_{s-} I^\varsigma F(r)] \\ & \leq [F(r) + F(s)] \beta_\varsigma(2, 2) + \frac{d^\varsigma(r, s)}{(s-r)^\varsigma} \end{aligned}$$

Corollary 2.4.

If $h(\ell) = 1^\varsigma$ in above inequality then following inequality holds

$$\begin{aligned} & \frac{1}{2^\varsigma h^\varsigma(\frac{1}{2})} F\left(\frac{r+s}{2}\right) - B_1 \leq \frac{1}{2^\varsigma (s-r)^\varsigma} [{}_{r+} I^\varsigma F(s) + {}_{s-} I^\varsigma F(r)] \\ & \leq \frac{[F(r) + F(s)]}{\Gamma(1+\varsigma)} + \frac{d^\varsigma(r, s)}{(s-r)^\varsigma} \end{aligned}$$

Corollary 2.5.

If $h(\ell) = \ell^{s\varsigma}$ in above inequality then following inequality holds

$$\begin{aligned} \frac{1}{2^\varsigma h^\varsigma(\frac{1}{2})} F\left(\frac{r+s}{2}\right) - B_1 &\leq \frac{1}{2^\varsigma (s-r)^\varsigma} [{}_r I^\varsigma F(s) + {}_s I^\varsigma F(r)] \\ &\leq [F(r) + F(s)] \frac{\Gamma(1+s\varsigma)}{\Gamma(1+(1+s)\varsigma)} + \frac{d^\varsigma(r,s)}{(s-r)^\varsigma} \end{aligned}$$

Theorem 2.2.

Let $F : [r, s] \rightarrow \mathbb{R}^\varsigma$ and $F(\cdot) \in I^\varsigma[r, s]$ be an approximately h -convex on $[r, s]$, with $0 \leq r < s$. Then following generalized local fractional integrals inequalities hold:

$$\begin{aligned} \frac{1}{2^\varsigma h^\varsigma(\frac{1}{2})} F\left(\frac{r+s}{2}\right) - B_2 &\leq \frac{1}{2^\varsigma \Lambda^\varsigma(1)} [{}_r I_\varphi^\varsigma F(s) + {}_s I_\varphi^\varsigma F(r)] \\ &\leq \frac{[F(r) + F(s)]}{2^\varsigma \Lambda^\varsigma(1)} \frac{1}{\Gamma(1+s\varsigma)} \int_0^1 \frac{\varphi_\varsigma((s-r)\ell)^\varsigma}{\ell^\varsigma} [h^\varsigma(\frac{\ell}{2}) + h^\varsigma(\frac{2-\ell}{2})] (d\ell)^\varsigma + d^\varsigma(r,s) \end{aligned}$$

where,

$$B_2 := \frac{1}{2^\varsigma \Lambda^\varsigma(1) h^\varsigma(\frac{1}{2}) \Gamma(1+s\varsigma)} \int_{\frac{r+s}{2}}^s \frac{\varphi_\varsigma(s-)^\varsigma}{(s-)^^\varsigma} d^\varsigma(\cdot, r+s-)(d)^\varsigma.$$

Proof. As F is generalized approximately h -convex function, so we have following inequality

$$F\left(\frac{u+v}{2}\right) \leq h^\varsigma\left(\frac{1}{2}\right) [F(u) + F(v)] + d^\varsigma(u, v) \quad (14)$$

Taking $u = \frac{\ell}{2}r + \frac{(2-\ell)}{2}s$ and $v = \frac{\ell}{2}s + \frac{(2-\ell)}{2}r$

$$\begin{aligned} \frac{1}{h^\varsigma(\frac{1}{2})} F\left(\frac{r+s}{2}\right) &\leq F\left(\frac{\ell}{2}r + \frac{(2-\ell)}{2}s\right) \\ &+ F\left(\frac{\ell}{2}s + \frac{(2-\ell)}{2}r\right) + \frac{1}{h^\varsigma(\frac{1}{2})} d\left(\frac{\ell}{2}r + \frac{(2-\ell)}{2}s, \frac{\ell}{2}s + \frac{(2-\ell)}{2}r\right) \end{aligned}$$

Multiply both sides of the inequality by $\frac{\varphi_\varsigma((s-r)\ell)}{\ell}$ and integrating local fractionally with respect to ℓ over $[0, 1]$, we obtain

$$\begin{aligned} &\frac{1}{h^\varsigma(\frac{1}{2})} F\left(\frac{r+s}{2}\right) \frac{1}{\Gamma(1+s\varsigma)} \int_0^1 \frac{\varphi_\varsigma((s-r)\ell)}{\ell} (d\ell)^\varsigma \\ &\leq \frac{1}{\Gamma(1+s\varsigma)} \int_0^1 \frac{\varphi_\varsigma((s-r)\ell)}{\ell} F\left(\frac{\ell}{2}r + \frac{(2-\ell)}{2}s\right) (d\ell)^\varsigma + \frac{1}{\Gamma(1+s\varsigma)} \int_0^1 \frac{\varphi_\varsigma((s-r)\ell)}{\ell} F\left(\frac{\ell}{2}s + \frac{(2-\ell)}{2}r\right) (d\ell)^\varsigma \\ &+ \frac{1}{h^\varsigma(\frac{1}{2}) \Gamma(1+s\varsigma)} \int_0^1 \frac{\varphi_\varsigma((s-r)\ell)}{\ell} d\left(\frac{\ell}{2}r + \frac{(2-\ell)}{2}s, \frac{\ell}{2}s + \frac{(2-\ell)}{2}r\right) (d\ell)^\varsigma \end{aligned}$$

As a consequence, we get

$$\begin{aligned} \frac{1}{2^\varsigma h^\varsigma(\frac{1}{2})} F\left(\frac{r+s}{2}\right) &\leq \frac{1}{2^\varsigma \Lambda^\varsigma(1)} [{}_{(\frac{r+s}{2})} I_\varphi^\varsigma F(s) + {}_{(\frac{r+s}{2})} I_\varphi^\varsigma F(r)] \\ &+ \frac{1}{2^\varsigma \Lambda^\varsigma(1) h^\varsigma(\frac{1}{2}) \Gamma(1+s\varsigma)} \int_{(\frac{r+s}{2})}^s \frac{\varphi_\varsigma(s-)^^\varsigma}{(s-)^^\varsigma} d^\varsigma(\cdot, r+s-)(d)^\varsigma. \end{aligned}$$

we proved the right hand side of the inequality.

To prove left hand side of the inequality, since F is approximately h -convex, we have

$$F\left(\frac{\ell}{2}r + \frac{(2-\ell)}{2}s\right) + F\left(\frac{\ell}{2}s + \frac{(2-\ell)}{2}r\right) \leq [F(r) + F(s)] \left[h^\varsigma \left(\frac{\ell}{2}\right) + h^\varsigma \left(\frac{2-\ell}{2}\right) \right] + 2^\varsigma d^\varsigma(r, s)$$

Multiplying both sides of inequality by $\frac{\varphi((s-r)\ell)}{\ell}$ and integrating local fractionally with respect to ℓ over $[0, 1]$, we obtain

$$\begin{aligned} & \frac{1}{\Gamma(1+\varsigma)} \int_0^1 \frac{\varphi((s-r)\ell)}{\ell} F\left(\frac{\ell}{2}r + \frac{(2-\ell)}{2}s\right) (d\ell)^\varsigma \\ & + \frac{1}{\Gamma(1+\varsigma)} \int_0^1 \frac{\varphi((s-r)\ell)}{\ell} F\left(\frac{\ell}{2}s + \frac{(2-\ell)}{2}r\right) (d\ell)^\varsigma + 2^\varsigma d^\varsigma(r, s) \frac{1}{\Gamma(1+\varsigma)} \int_0^1 \frac{\varphi((s-r)\ell)}{\ell} (d\ell)^\varsigma \\ & \leq [F(r) + F(s)] \frac{1}{\Gamma(1+\varsigma)} \int_0^1 \frac{\varphi((s-r)\ell)}{\ell} \left[h^\varsigma \left(\frac{\ell}{2}\right) + h^\varsigma \left(\frac{2-\ell}{2}\right) \right] (d\ell)^\varsigma + 2^\varsigma d^\varsigma(r, s) \Lambda^\varsigma(1) \end{aligned}$$

As a result, we get

$$\begin{aligned} & \frac{1}{2^\varsigma \Lambda^\varsigma(1)} [({}_{(r+\frac{s}{2})+}I_\varphi^\varsigma F(s) + ({}_{(r+\frac{s}{2})-}I_\varphi^\varsigma F(r))] \\ & \leq \frac{[F(r) + F(s)]}{2^\varsigma \Lambda^\varsigma(1) \Gamma(1+\varsigma)} \int_0^1 \frac{\varphi((s-r)\ell)}{\ell} \left[h^\varsigma \left(\frac{\ell}{2}\right) + h^\varsigma \left(\frac{2-\ell}{2}\right) \right] (d\ell)^\varsigma + d^\varsigma(r, s) \end{aligned}$$

□

Corollary 2.6.

If $\varphi(\ell) = \ell^\varsigma$ then following inequality holds

$$\begin{aligned} & \frac{1}{2^\varsigma h^\varsigma(\frac{1}{2})} F\left(\frac{r+s}{2}\right) - B_2 \leq \frac{1}{2^\varsigma (s-r)^\varsigma} [({}_{(r+\frac{s}{2})+}I^\varsigma F(s) + ({}_{(r+\frac{s}{2})-}I^\varsigma F(r))] \\ & \leq \frac{[F(r) + F(s)]}{2^\varsigma} \frac{1}{\Gamma(1+\varsigma)} \int_0^1 \left[h^\varsigma \left(\frac{\ell}{2}\right) + h^\varsigma \left(\frac{2-\ell}{2}\right) \right] (dt)^\varsigma + \frac{d^\varsigma(r, s)}{(s-r)^\varsigma} \end{aligned}$$

where,

$$B_2 := \frac{1}{2^\varsigma h^\varsigma(\frac{1}{2}) \Gamma(1+\varsigma) (s-r)^\varsigma} \int_{(r+\frac{s}{2})}^s d^\varsigma(r, s-)(d)^\varsigma.$$

Corollary 2.7.

If $h(\ell) = \ell^\varsigma$ in above inequality then following inequality holds

$$\begin{aligned} & \frac{1}{2^\varsigma h^\varsigma(\frac{1}{2})} F\left(\frac{r+s}{2}\right) - B_2 \leq \frac{1}{2^\varsigma (s-r)^\varsigma} [({}_{(r+\frac{s}{2})+}I^\varsigma F(s) + ({}_{(r+\frac{s}{2})-}I^\varsigma F(r))] \\ & \leq \frac{[F(r) + F(s)]}{\Gamma(1+\varsigma)} + \frac{d^\varsigma(r, s)}{(s-r)^\varsigma} \end{aligned}$$

Corollary 2.8.

If $h(\ell) = \ell^\varsigma(1-\ell)^\varsigma$ in above inequality then following inequality holds

$$\begin{aligned} & \frac{1}{2^\varsigma h^\varsigma(\frac{1}{2})} F\left(\frac{r+s}{2}\right) - B_2 \leq \frac{1}{2^\varsigma (s-r)^\varsigma} [({}_{(r+\frac{s}{2})+}I^\varsigma F(s) + ({}_{(r+\frac{s}{2})-}I^\varsigma F(r))] \\ & \leq [F(r) + F(s)] \left[\frac{\beta_\varsigma(2, 2)}{2^\varsigma} + \frac{\Gamma(1+\varsigma)}{4^\varsigma \Gamma(1+\varsigma)} - \frac{\Gamma(1+2\varsigma)}{8^\varsigma \Gamma(1+3\varsigma)} \right] + \frac{d^\varsigma(r, s)}{(s-r)^\varsigma} \end{aligned}$$

Corollary 2.9.

If $h(\ell) = 1^\varsigma$ in above inequality then following inequality holds

$$\begin{aligned} \frac{1}{2^\varsigma h^\varsigma(\frac{1}{2})} F\left(\frac{r+s}{2}\right) - B_2 &\leq \frac{1}{2^\varsigma (s-r)^\varsigma} [({}_{(r+\frac{s}{2})+}I^\varsigma F(s) + ({}_{(r+\frac{s}{2})-}I^\varsigma F(r))] \\ &\leq \frac{[F(r) + F(s)]}{\Gamma(1+\varsigma)} + \frac{d^\varsigma(r, s)}{(s-r)^\varsigma} \end{aligned}$$

Corollary 2.10.

If $h(\ell) = \ell^{s\varsigma}$ in above inequality then following inequality holds

$$\begin{aligned} \frac{1}{2^\varsigma h^\varsigma(\frac{1}{2})} F\left(\frac{r+s}{2}\right) - B_2 &\leq \frac{1}{2^\varsigma (s-r)^\varsigma} [({}_{(r+\frac{s}{2})+}I^\varsigma F(s) + ({}_{(r+\frac{s}{2})-}I^\varsigma F(r))] \\ &\leq \frac{[F(r) + F(s)]}{\Gamma(1+\varsigma)} + \frac{d^\varsigma(r, s)}{(s-r)^\varsigma} \end{aligned}$$

3. Conclusion

This study presented the concept of generalized approximately h -convex function on fractal sets. Involving generalized local fractional integral operators, we proved some latest generalized local fractional integral inequalities. Some special cases of novel findings are also established. It is to be expected that the the notion introduced in this study may open new doors for researchers to work on generalized approximately h -convex function in different directions.

References

-
- [1] A. Atangana, Modelling the spread of COVID-19 with new fractal-fractional operators: Can the lockdown save mankind before vaccination?. *Chaos, Solitons and Fractals*, **136** (2020) 109860.
 - [2] O. Almutairi, A. Kilicman, Generalized Fejér–Hermite–Hadamard type via generalized $(h - m)$ -convexity on fractal sets and applications. *Chaos, Solitons & Fractals*, **147**(2021) 110938.
 - [3] M.W. Alomari, The Hermite-Hadamard inequality on hypercuboid.(2016) arXiv preprint arXiv:1604.01857.
 - [4] M. Bibi, and M. Muddassar, Hermite-Hadamard type fractional integral inequalities for strongly generalized-prequasi-invex function. *International Journal of Nonlinear Analysis and Applications*.(2022)
 - [5] S. I. Butt, A. Kashuri, M. Tariq, J. Nasir, A. Aslam and W. Gao, Hermite–Hadamard-type inequalities via n -polynomial exponential-type convexity and their applications. *Advances in Difference Equations*, **1**(2020) 1-25.
 - [6] H. H. Chu, S. Rashid, Z. Hammouch and Y. M. Chu, New fractional estimates for Hermite-Hadamard-Mercer’s type inequalities. *Alexandria Engineering Journal*, **59**(2020) no. 5 3079-3089.
 - [7] T. Du, M. U. Awan, A. Kashuri and S. Zhao, Some k -fractional extensions of the trapezium inequalities through generalized relative semi- (m, h) -preinvexity. *Applicable Analysis* (2019) 1-21.

-
- [8] T. Du, H. Wang, M. A. Khan and Y. Zhang, Certain integral inequalities considering generalized m -convexity on fractal sets and their applications. *Fractals*, **27**(2019) no.7, 1950117.
- [9] J. Hadamard, Étude sur les propriétés des fonctions entières et en particulier d'une fonction considérée par Riemann. *Journal de mathématiques pures et appliquées*.(1893), 171-216.
- [10] S. Iftikhar, S. Erden, P. Kumam and M.U.Awan, Local fractional Newton's inequalities involving generalized harmonic convex functions. *Advances in Difference Equations*, **2020**(2020) 1-14.
- [11] A. Kashuri, H. Budak, R. Kapucu and R. Liko, New Inequalities for approximately h -convex functions pertaining generalized fractional integrals.DOI: 10.37560/matbil21452151k
- [12] A. Kashuri, M. Raees, and M. Anwar, Some integral inequalities for approximately h -convex functions and their applications. *Proyecciones (Antofagasta)*, **40** (2021) no. 2 , pp.481-504.
- [13] S. Kumar, A. Atangana, A numerical study of the nonlinear fractional mathematical model of tumor cells in presence of chemotherapeutic treatment. *International Journal of Biomathematics*, **13**(2020) no. 3, 2050021.
- [14] H. Mo, X. Sui D.Yu, Generalized convex functions on fractal sets and two related inequalities. In *Abstract and Applied Analysis* 2014(2014, January) Hindawi.
- [15] M. Muddassar, S. Dragomir, Z. Hussain, Rayna's fractional integral operations on Hermite–Hadamard inequalities with η - G -Preinvex functions. *Advances in Operator Theory*,(2020), 1-16.
- [16] M. Muddassar, and F. Bibi, 2022. A Short note on the fractional trapezium type integral inequalities. *Innovative Journal of Mathematics (IJM)*, **1**(2022), no. 2,63-70.
- [17] M. Z. Sarikaya, F. Ertuğral, On the generalized Hermite-Hadamard inequalities, *Ann. Univ. Craiova Math. Comput. Sci. Ser.* **47**(2020), 193-213.
- [18] M. Sarikaya, H. Budak, Generalized Ostrowski type inequalities for local fractional integrals, *Proceedings of the American Mathematical Society*, **145**(2017) no. 4, 1527-1538.
- [19] Y. Q. Song, S.I. Butt, A. Kashuri, J. Nasir and M. Nadeem, New fractional integral inequalities pertaining 2D-approximately coordinate $(r_1, h_1) - (r_2, h_2)$ -convex functions. *Alexandria Engineering Journal*, **61**(2022) no. 1, 563-573.
- [20] W. Sun, Hermite-Hadamard type local fractional integral inequalities for generalized s -preinvex functions and their generalization. *Fractals*, **29** (2021) no.4
- [21] W. Sun, Generalized h -convexity on fractal sets and some generalized Hadamard-Type inequalities, *Fractals*, **28** (2020) no.2, 2050021.
- [22] W. Sun, Some Hermite–Hadamard type inequalities for generalized h -preinvex function via local fractional integrals and their applications. *Advances in Difference Equations*, **2020**(2020) no.1, 1-14.
- [23] W. Sun, Some local fractional integral inequalities for generalized preinvex functions and applications to numerical quadrature, *Fractals*, **27** (2019) no.05, 1950071.
- [24] W.Sun, Some new inequalities for generalized h -convex functions involving local fractional integral operators with Mittag-Leffler kernel. *Mathematical Methods in the Applied Sciences* **44** (2021) no.6, 4985-4998.

- [25] W. Sun, Hermite-Hadamard type local fractional integral inequalities with Mittag-Leffler kernel for generalized preinvex functions *Fractals* (2021).<https://doi.org/10.1142/S0218348X21502534>
- [26] S. Varoanec, On h -convexity, *J. Math. Anal. Appl.*, **326** (2007), 303-311.
- [27] K. J. Wang, Variational principle and approximate solution for the generalized Burgers-Huxley equation with fractal derivative, *Fractals*, **29** (2021), no. 2, 2150044-1246.
- [28] X. J. Yang, *Advanced local fractional calculus and its applications*, 2012.
- [29] X. J. Yang, D. Baleanu, H. M. Srivastava, *Local fractional integral transforms and their applications*, ACADEMIC PRESS, 2015
- [30] X. J. Yang, F. Gao and H. M. Srivastava, Exact travelling wave solutions for the local fractional two-dimensional Burgers-type equations. *Computers & Mathematics with Applications*, **73** (2017) no.2, 203-210.